

Solution of classical stochastic one dimensional many-body systems

P.-A. Bares and M. Mobilia

Institut de Physique Théorique, École Polytechnique Fédérale de Lausanne, CH-1015 Lausanne
(July , 1999)

We propose a simple method that allows, in one dimension, to solve exactly a wide class of classical stochastic many-body systems far from equilibrium. For the sake of illustration and without loss of generality, we focus on a model that describes the asymmetric diffusion of hard core particles in the presence of an external source and instantaneous annihilation. Starting from a Master equation formulation of the problem we show that the density and multi-point correlation functions obey a closed set of integro-differential equations which in turn can be solved numerically and/or analytically.

Recently, nonequilibrium statistical mechanics in low dimensional systems has attracted much interest [1]. Whereas classical statistical mechanics has a well established body of concepts and tools to deal with systems at or near thermodynamic equilibrium, the situation is less satisfactory for far from equilibrium systems. The theoretical understanding of classical stochastic many-body systems where both purely diffusive motion (due for example to static disorder, phonons, etc.) and reactions (for example fusion, annihilation, coagulation, etc., where the number of particles is not conserved) take place simultaneously is relevant to a wide class of phenomena in physics, biology, economics, a.s.o.

In view of this, we have chosen to investigate an important class of models describing diffusion-limited reactions (for recent reviews see [2]). While it has long been recognized that the mean field approach is not applicable in one and two dimensions, it is only recently that much progress has been achieved in the description of the anomalous kinetics of low dimensional systems. Some open problems in this field are reminiscent of those studied intensively in the field of strongly correlated low-dimensional systems. Hence, one might hope to use with advantage the powerful tools devised in the latter.

The last decade has witnessed new developments in the Master equation approach. The Markov generator of the stochastic time evolution for the probability distribution can be interpreted as the Hamiltonian of quantum spin chains (with quantum group symmetries) or alternatively of fermion or boson many-body systems with interaction terms. The dynamics is then coded in an imaginary time Schrödinger equation or alternatively formulated as an Euclidean field theory. The most formal approach expresses the Hamiltonian as a sum of generators of an Hecke Algebra and, through Baxterization, associates an integrable vertex model [3,4]. The free fermion models considered by Grynberg et al. [5], Schütz and others (see [2] and references therein) can be viewed as special cases of the above with a grading of the Yang-Baxter algebra. As is known in the field of integrable models, it is in

general extremely difficult, if not impossible, to extract the correlations from the Bethe “wave functions”, even in their algebraic form (see for example [6]). This is unfortunate because conceptually as well as experimentally (see [1] and references therein), the correlations coded in these “wave functions” are the most relevant quantities. We should mention at this point however that there is one successful application of ideas developed in the field of strongly correlated systems, namely the extension of the numerical Density Matrix Renormalization Group to nonequilibrium statistical mechanics in one dimension [7].

On the analytical front, a powerful method is that developed by Cardy and collaborators [8]. Assuming that the system is in a low-density regime, these authors rewrite the Markov generator of the stochastic time evolution as a bosonic Hamiltonian which in turn can be treated by field theory and renormalization group techniques in arbitrary dimension. For certain models, it is even possible to evaluate exactly the beta functions for the coupling constants [8]. This is a first indication that the underlying mathematical structure is substantially simpler than that of the quantum many-body problem. To our knowledge, this observation as well as its implications do not seem to have drawn much attention in the literature.

Despite the progress achieved in the last decade, there are still many open or poorly understood problems, even in one dimension: the multi-species and the epidemic models, the propagation of catalytic shock waves, etc. In response to these challenges, we were lead to consider a fermionic field theory formulation of the Master equation for dense systems [9]. While inquiring about the possible application of the renormalization group to the fermion field theory, we realized that it is possible to express the density and correlation functions as solutions of a closed set of integro-differential equations. These in turn can be solved exactly numerically and/or analytically. This is an unexpected and remarkable result, especially in view of the large amount of work that has been performed in the field using sophisticated methods. Our approach in contrast is simple and elementary. In this letter we would like to report on this rather unanticipated observation.

For illustrative purposes, we focus here on an extended Lushnikov model [10] that describes the (asymmetric) diffusion of particles in the presence of bilinear source and annihilation. Certainly, such a model does not constitute a prototype for classical stochastic many-body systems as does the Ising model for phase transitions: the variety of phenomena to be described in nonequilibrium statistical mechanics is much too rich to be coded in a single model. Though we are perfectly aware of the numerous potential applications that our method provides, we will refrain from mentioning them here due to a lack of space and refer the reader to elsewhere (e.g. [1,2]).

Consider a lattice of L ($L/2$ even) sites with periodic boundary conditions on which classical particles with

hard core can diffuse with rates h to the left and h' to the right, respectively. On encounter, pairs of particles annihilate with rate ϵ' . A source injects particle pairs with rate ϵ . Rewriting the Master equation as a Schrödinger equation in imaginary time, we have $d|P(t)\rangle/dt = -H|P(t)\rangle$, with $(\sigma_{m+L}^+ = \sigma_m^+)$

$$H = - \sum_{m=1}^L \left\{ \begin{aligned} & h\sigma_{m+1}^+ \sigma_m^- + h'\sigma_m^+ \sigma_{m+1}^- + \epsilon'\sigma_m^- \sigma_{m+1}^- + \epsilon(\sigma_m^+ \sigma_{m+1}^+ - 1) \\ & + (2\epsilon - h - h')\sigma_m^+ \sigma_m^- - (\epsilon' + \epsilon - h - h')\sigma_m^+ \sigma_m^- \sigma_{m+1}^+ \sigma_{m+1}^- \end{aligned} \right\} \quad (1)$$

where $\sigma_m^\pm$ denote the standard Pauli matrices. The Dirac ket $|P(t)\rangle = \sum_\eta P_\eta(t)|\eta\rangle$, where η denotes a random configuration in Fock space, encodes the probabilities of the various configurations (see [5]). The above dynamics is supplemented with the initial conditions $|P(0)\rangle = |\rho\rangle$ (see below). It is convenient to introduce the left vacuum $\langle\tilde{\chi}| = \sum_\eta \langle\eta|$. Conservation of probability implies then $\langle\tilde{\chi}|H = 0$, while expectation values are linear in the kets $\langle A(t)\rangle = \langle\tilde{\chi}|A|P(t)\rangle$. The models that have been solved previously correspond to $\epsilon' + \epsilon = h + h'$ (see e.g. [2,5,9,10]), i.e., to quadratic hamiltonians. Below, we show that the Hamiltonian H can be solved exactly for $\gamma = \epsilon' + \epsilon - (h + h') \neq 0$ by elementary means, i.e., that the density and two-point (multi-point) correlation function(s) can be computed exactly even though the hamiltonian is non-quadratic.

Performing a Jordan-Wigner transformation, and subsequently a Fourier transformation, leads to

$$H = \sum_{q>0} \left[\omega(q)a_q^\dagger a_q + \bar{\omega}(q)a_{-q}^\dagger a_{-q} + 2\sin q \left(\epsilon' a_q a_{-q} + \epsilon a_{-q}^\dagger a_q^\dagger \right) \right] + \epsilon L \\ - \gamma \sum_q (1 - \cos q) a_q^\dagger a_q + \frac{\gamma}{L} \sum_{q_1, q_2, q_3} \cos(q_1 - q_2) a_{q_1}^\dagger a_{q_2} a_{q_3}^\dagger a_{q_1+q_3-q_2}, \quad (2)$$

where $\omega(q) = a - b \cos q + i(h' - h) \sin q$, with $a = \epsilon' - \epsilon$, $b = \epsilon' + \epsilon$. Following reference [2], we consider a translationally invariant and uncorrelated initial state with an even number N of particles of density $\rho = N/L$, i.e.,

$$|\rho\rangle = \frac{2(1-\rho)^L}{1+(1-2\rho)^L} \prod_{q>0} \left(1 - \mu^2 \cot\left(\frac{q}{2}\right) a_{-q}^\dagger a_q^\dagger \right) |0\rangle, \quad (3)$$

where $\mu = \rho/(1-\rho)$.

Using the properties of the left vacuum ($\langle\tilde{\chi}|a_q^\dagger = \cot(\frac{q}{2})\langle\tilde{\chi}|a_{-q}$, see [2]), and with the notation $\langle a_{-q} a_{q'} \rangle(t) \equiv \langle\tilde{\chi}|a_{-q} a_{q'} e^{-Ht}|\rho\rangle$, the density and the two-point density correlation function can be written as

$$\rho(t) = \langle a_m^\dagger a_m \rangle(t) = \frac{1}{L} \sum_{q, q'} e^{i(q-q')m} \cot\left(\frac{q}{2}\right) \langle\tilde{\chi}|a_{-q} a_{q'} e^{-Ht}|\rho\rangle \quad (4)$$

$$\begin{aligned} \langle n_m n_n \rangle(t) = \frac{1}{L^2} \sum_{k, k', q, q'} e^{im(q-q') + in(k-k')} & \left[\langle a_{-q} a_{q'} \rangle(t) \delta_{k', k} - \langle a_{q'} a_{k'} \rangle(t) \delta_{q, -k} \right. \\ & \left. + \cot\left(\frac{k}{2}\right) \langle a_{-k} a_{-q} a_{q'} a_{k'} \rangle(t) \right] \cot\left(\frac{q}{2}\right) \quad (5) \end{aligned}$$

Futhermore, it can be shown that [11]

$$\langle \tilde{\chi} | a_{-q} a_{q'} e^{-Ht} | \rho \rangle = g(q, t) \langle a_{-q} a_{q'} \rangle(0) = g(q, t) \frac{\mu^2 \cot(\frac{q}{2})}{1 + \mu^2 \cot(\frac{q}{2})^2} \delta_{q, q'}, \quad (6)$$

where $g(q, t)$ is a function (even in q) that is a solution of a non-linear integro-differential equation (see below), with $g(q, t = 0) = 1$, $g(q, t < 0) = 0$ and $g(q, t) = g(q + 2\pi, t)$. This remarkable property can be generalized to higher order correlation functions. Due to the conservation of probability, an analog of the Wick theorem applies and all multipoint correlation functions can be computed. For simplicity's sake we consider in the following a lattice initially full, i.e., $\rho = 1$ (for the general case see [11]) and obtain the integro-differential equation for $g(q, t)$

$$\begin{aligned} \frac{\partial}{\partial t} g(q, t) = 2\epsilon(1 + \cos q) - 2(b - \gamma)g(q, t) + 2(a - \gamma) \cos q g(q, t) \\ - \frac{\gamma}{\pi}(1 + \cos q) \int_{-\pi}^{\pi} dk (1 - \cos k) g(k, t) - \frac{2\gamma}{\pi} g(q, t) \int_{-\pi}^{\pi} dk \cos k g(k, t) \\ + \frac{2\gamma}{\pi} g(q, t) \cos q \int_{-\pi}^{\pi} dk g(k, t) \end{aligned} \quad (7)$$

with the initial condition $g(q, 0) = 1$. This equation with the appropriate boundary conditions fully solves the problem of the time evolution of the density and correlation function. Indeed, introducing the help functions $F_n(t) = (1/2\pi) \int_{-\pi}^{\pi} dq g(q, t) \cos(nq)$ (for n integer), the density can be written as

$$\rho(t) = F_0(t), \quad (8)$$

while it is convenient to introduce the function $\mathcal{G}(r, t) \equiv 1 - 2 \sum_{1 \leq n \leq r} F_{2n-r-1}(t)$ to express the (connected) two-point correlation function as

$$\mathcal{C}_r(t) = \frac{1}{4} [\mathcal{G}(r-1, t) \mathcal{G}(r+1, t) - \mathcal{G}^2(r, t)] \quad (9)$$

The help functions $F_n(t)$ have the integral representation

$$\begin{aligned} F_n(t) = e^{-\tilde{K}_1(t)} I_n(\tilde{K}_0(t)) \\ + \int_0^t d\tau \left(\epsilon + \gamma \{F_1(t-\tau) - F_0(t-\tau)\} \right) e^{\tilde{K}_1(t-\tau) - \tilde{K}_1(t)} \left\{ 2I_n(\tilde{K}_0(t) - \tilde{K}_0(t-\tau)) \right. \\ \left. + I_{n+1}(\tilde{K}_0(t) - \tilde{K}_0(t-\tau)) + I_{n-1}(\tilde{K}_0(t) - \tilde{K}_0(t-\tau)) \right\}, \end{aligned} \quad (10)$$

where $I_n(z)$ denote the modified Bessel functions of imaginary argument, and $\tilde{K}_0(t) \equiv 2(a - \gamma)t + 4\gamma \int_0^t ds F_0(s)$, $\tilde{K}_1(t) \equiv 2(b - \gamma)t + 4\gamma \int_0^t ds F_1(s)$. It

is worth noting that $F_0(t)$ and $F_1(t)$ are obtained self-consistently while the higher order functions $F_n(t)$ ($n \geq 2$) follow from simple integration. Notice that at $\gamma = 0$, we recover the results obtained in [2,4,5,9,10]. We have solved the integral equations (10) and obtained the $F_n(t)$ numerically as well as analytically in the asymptotic regime. In the long time regime, the asymptotes follow as

i) In the absence of source ($ht, h't, \epsilon't \gg 1, \epsilon = 0$)

$$\rho(t) \approx \frac{1}{\sqrt{4\pi(h+h')t}} - \frac{\gamma}{\pi\epsilon'(h+h')t} \quad (11)$$

$$\mathcal{C}_r(t) \approx \frac{-1}{4\pi(h+h')t} + \left\{ \pi r + \frac{8\gamma}{\epsilon'} \right\} \frac{1}{(4\pi(h+h')t)^{3/2}} \quad (12)$$

The result for the density agrees with that of reference [12]. The numerical computation of $\rho(t)$ is shown in Fig.1.

ii) In the presence of source ($ht, h't, |a|t, bt \gg 1$), two regimes have to be distinguished, namely

1) If $\tilde{K}_0(t) > 0$, then,

$$\rho(t) - \rho(\infty) \sim -\mathcal{C}_r(t) \sim \frac{\exp(-4(\epsilon - \gamma\rho(\infty)^2)t)}{t^{3/2}} \quad (13)$$

A numerical illustration of this case is shown in Fig.2

2) If $\tilde{K}_0(t) < 0$, then,

$$\rho(t) - \rho(\infty) \sim \mathcal{C}_r(t) \sim \frac{\exp\left(-2[(b-\gamma) + 2\gamma F_1(\infty) - |a-\gamma + 2\gamma\rho(\infty)|]t\right)}{t^{1/2}} \quad (14)$$

where $\rho(\infty) = 1/(1 + \sqrt{\epsilon'/\epsilon})$ is the stationary density which coincides with the (free fermion) $\gamma = 0$ case, and $F_1(\infty) = \sqrt{\epsilon\epsilon'}/(\sqrt{\epsilon} + \sqrt{\epsilon'})^2$.

The results (11)-(14) indicate that the γ term is irrelevant in the long time limit in the sense that it does not renormalize the exponent of the algebraic (in the absence of source) or of the subdominant-algebraic (in the presence of source) decay. Such a behavior had been observed numerically (in the absence of source) in reference [12]. The correlation times are however renormalized.

The extension of our method to deal with epidemic models and many species is indeed possible. Local sources or sinks as well as boundary (initial state) terms associated with local operators in the hamiltonian formulation can also be dealt with whenever the Jordan-Wigner phase can be made to cancel. A physical illustration of this is realized in a model of photoluminescent diodes with injection currents at the boundaries. These and other physically interesting applications of the method proposed in this paper will be discussed elsewhere [11].

ACKNOWLEDGMENTS

We thank J. Cardy, B. Douçot, M. Droz, Ch. Gruber, Y. Hatsugai, Ph. Nozières, T.M. Rice, G.M. Schütz, A. Tsvelik and X.-G. Wen for discussions. We are especially grateful to K. Grzegorzczuk, S.Gyger and L.Klinger for computational assistance. The support of the Swiss National Fonds is gratefully acknowledged.

-
- [1] *Nonequilibrium Statistical mechanics in One dimension*, ed. by V. Privman, Cambridge University Press, 1997.
 - [2] J.E. Santos, G.M. Schütz, R.B. Stinchcombe, J. Chem. Phys. **105**, 2399, (1996); D. C. Mattis and M. L. Glasser, Rev. Mod. Phys. **70**, 979 (1998); G.M. Schütz, *Integrable Stochastic Many-Body Systems*, to appear in *Phase Transition and critical Phenomena*, Domb and Lebowitz eds. (1999)
 - [3] V.R. Jones, Int. J. Mod. Phys.B **4**, 701 (1990)
 - [4] F.C. Alcaraz, M. Droz, M. Henkel and V. Rittenberg, Annals of Physics **230**, 250-302 (1994)
 - [5] M.D. Grynberg and R.B. Stinchcombe, Phys. Rev. Lett. **74**, 1242 (1995); M.D. Grynberg and R.B. Stinchcombe, Phys. Rev. Lett. **76**, 851 (1996)
 - [6] V.E. Korepin, N.M. Bogoliubov and A.G. Izergin, *Quantum Inverse Scattering Method and Correlation Functions*, Cambridge University press, 1993.
 - [7] E. Carlon, M. Henkel, U. Schwoillwoeck, e-print cond-mat/9902041
 - [8] B.P. Lee and J.L. Cardy, J. Stat. Phys. **80**, 971 (1995); J. Cardy, cond-mat/9607163v2, September 1996; J.L. Cardy and U.C. Täuber, J.Stat.Phys. **90**, 1 (1998); J.L. Cardy, Cours de Troisième Cycle de Suisse Romande, *Field Theory and Nonequilibrium Statistical Mechanics*, April 1999
 - [9] P.-A. Bares and M. Mobilia, Phys. Rev. E **59**, 1996 (1999)
 - [10] A.A. Lushnikov, Sov. Phys. JETP **64**(4), 811 (1986)
 - [11] P.-A. Bares and M. Mobilia, in preparation.
 - [12] K. Krebs, M.P. Pfannmüller, H. Simon and B. Wehefritz, J. Stat. Phys. **78**, 1471 (1995).




FIG. 1. Density $\rho(t)$ versus time for $\epsilon = 0, \epsilon' = 1.5$ and $\gamma = 0.5$; (upper curve) $\gamma = 0$ (middle curve: free-fermions); $\gamma = -0.5$ (lower curve).




FIG. 2. The logarithm of the density $t^{3/2}[\rho(t) - \rho(\infty)]$ (stars, $\rho(\infty) = 1/3$) and of the two-point correlation function $-t^{3/2}C_r(t)$, $r=1$ (circles), $r=2$ (triangles) for $\epsilon = 0.5, \epsilon' = 2$, and $\gamma = 1$ versus time. We have included the shifted asymptote $\text{const} - 14t/9$ as a guide to the eye (thin line).